*IRINOTECAN HYDROCHLORIDE INJECTION

20 mg/mL of Irinotecan Hydrochloride (as the trihydrate)

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of administration	Dosage form/Strength	Clinically Relevant Nonmedicinal Ingredients
Intravenous	Solution / 20 mg/mL	*Sorbitol, for a complete listing see Dosage Forms, Composition and Packaging sections*

INDICATIONS AND CLINICAL USE

IRINOTECAN HYDROCHLORIDE INJECTION (irinotecan hydrochloride) is indicated as a component of first-line therapy for patients with metastatic carcinoma of the colon or rectum.

IRINOTECAN HYDROCHLORIDE INJECTION is indicated for the treatment of patients with metastatic carcinoma of the colon or rectum whose disease has recurred or progressed following 5-fluorouracil based therapy.

IRINOTECAN HYDROCHLORIDE INJECTION should be administered only under the supervision of a physician who is experienced in the use of cancer chemotherapeutic agents. Appropriate management of complications is possible only when adequate diagnostic and treatment facilities are readily available.

Geriatrics (≥ 65 years of age): Pharmacokinetic studies suggests that patients 65 years of age or older should be closely monitored because of a greater risk of severe toxicity. No specific dosing recommendations may apply to this population depending upon the regimen used (see WARNINGS AND PRECAUTIONS, Geriatrics, DOSAGE AND ADMINISTRATION, CLINICAL TRIALS).

Pediatrics: The safety and effectiveness of IRINOTECAN HYDROCHLORIDE INJECTION in the pediatric population has not been established (see WARNINGS AND PRECAUTIONS).

CONTRAINDICATIONS

IRINOTECAN HYDROCHLORIDE INJECTION is contraindicated in patients who are hypersensitive to this drug or to any ingredient in the formulation or component of the container.

For a complete listing, see the Dosage Forms, Composition and Packaging section of the product monograph.

Co-administration of IRINOTECAN HYDROCHLORIDE INJECTION with azole antifungals (ketoconazole, fluconazole, itraconazole), known CYP3A4 inhibitors, is contraindicated because this can lead to an increase in the relative exposure to the active metabolite SN-38 and can therefore possibly lead to increased toxicity.

Co-administration of IRINOTECAN HYDROCHLORIDE INJECTION with strong CYP3A4 inducers (anticonvulsants, e.g. phenytoin, carbamazepine, phenobarbital, primidone, or St John's Wort, rifampicin) is contraindicated because this can lead to a decrease in the relative exposure to the active metabolite SN-38 and can therefore possibly lead to decreased efficacy (see DRUG INTERACTIONS).

The appropriate starting dose of irinotecan when co-administered with CYP3A4 inhibitors has not been determined.

The appropriate starting dose of irinotecan when co-administered with CYP3A4 inducers has not been determined.

WARNINGS AND PRECAUTIONS

Serious Warnings and Precautions

- IRINOTECAN HYDROCHLORIDE INJECTION should be administered only under the supervision of a physician who is experienced in the use of cancer chemotherapeutic agents.
- Severe early and late forms of diarrhea can occur and require careful and electrolyte imbalance (See WARNINGS AND PRECAUTIONS, ADVERSE DRUG REACTIONS).
- Tightnitis, ulcerative and ischemic colitis, ileus and dehydration (See WARNINGS AND PRECAUTIONS, POST-MARKET ADVERSE DRUG REACTIONS).
- Severe myelosuppression with grade 3-4 neutropenia (See WARNINGS AND PRECAUTIONS, ADVERSE DRUG REACTIONS).

General

Patients at Particular Risk:

Clinical studies examine particular caution in monitoring the effects of IRINOTECAN HYDROCHLORIDE INJECTION in patients with poor performance status. Patients with performance status of 3 or 4 should not receive IRINOTECAN HYDROCHLORIDE INJECTION. In patients receiving either irinotecan/5-FU/LV or 5-FU/LV in clinical trials comparing these agents, higher rates of treatment discontinuation due to adverse events, treatment delays, dose reductions and treatment early deaths were observed in patients with a baseline performance status of 0 or 1. Close monitoring is recommended in patients who have previously received pelvic/abdominal irradiation or patients who may have reduced bone marrow tolerant of the toxic effects of the drug. The use of IRINOTECAN HYDROCHLORIDE INJECTION has not been established in patients with significant hepatic dysfunction (see WARNINGS AND PRECAUTIONS, Hepatic). There are known and suspected drug-drug interactions (see WARNINGS AND PRECAUTIONS, Drug Interactions and DRUG INTERACTIONS).

Drug Interactions:

CYP3A4 inhibitors:

- Co-administration of irinotecan with azole antifungals (ketoconazole, fluconazole, itraconazole) is contraindicated (see CONTRAINDICATIONS).
- Co-administration of irinotecan with other CYP3A4 inhibitors (eg., cimetidine, fluvoxaporine antibiotics [ciprofloxacin and norfloxacin in patients with compromised renal function], macrolide antibiotics [azithromycin, clarithromycin, erythromycin], protease inhibitors, grapefruit juice, and CYP3A4 inhibitory herbal products (echinacea) may increase the relative exposure to the active metabolite SN-38 and therefore possibly increase toxicity (see DRUG INTERACTIONS, DOSAGE AND ADMINISTRATION).

The appropriate starting dose of irinotecan when co-administered with CYP3A4 inhibitors has not been determined.

CYP3A4 Inducers:

- The co-administration of irinotecan with CYP3A4 inducers (eg. St. John's Wort, phenytoin, carbamazepine, rifampicin, phenobarbital, primidone) may result in a decrease in the active concentration of the active metabolite SN-38, which could possibly lead to a reduction in efficacy (see DRUG INTERACTIONS).

The appropriate starting dose of irinotecan when co-administered with CYP3A4 inducers has not been determined.

Irradiation Therapy:

The concurrent administration of IRINOTECAN HYDROCHLORIDE INJECTION with irradiation is not recommended.

Extravasation:

IRINOTECAN HYDROCHLORIDE INJECTION is administered by intravenous infusion. Care should be taken to avoid extravasation. The infusion site should be monitored for signs of inflammation or extravasation. If extravasation occurs, flushing with water and application of ice packs should be applied to the area are recommended.

Carcinogenesis and Mutagenesis

Carcinogenicity studies have not been conducted. Rats administered 2 mg/kg or 25 mg/kg irinotecan IV once weekly for 13 weeks and allowed to recover for 91 weeks had a significant linear trend with dose for the incidence of combined adenomas and carcinomas of small intestinal origin and endometrial stromal sarcomas. Irinotecan and SN-38 were not mutagenic in bacterial *in vitro* assays (Ames reverse mutation assay) or mammalian *in vitro* assays (mouse lymphoma L5178Y/TK+/- forward mutation assay) cells) and *in vivo* (micronucleus test in mice) (see TOXICOLOGY). However, irinotecan may be able to induce chromosomal damage in human spermatocytes. For this reason, males undergoing IRINOTECAN HYDROCHLORIDE INJECTION treatment should discuss effective contraceptive methods with their doctors.

Cardiovascular

All thromboembolic events (includes angina pectoris, arterial thrombosis, cerebral infarct, cerebrovascular accident, deep thrombophlebitis, embolism, myocardial infarct, myocardial ischemia, peripheral vascular disorder, pulmonary embolus, sudden death, thrombophlebitis, venous thrombosis) were reported in 4.8% of patients receiving irinotecan. The most commonly observed in patients receiving irinotecan. However, a definite cause of these events has not been determined. See also **Clinical Trial Adverse Drug Reactions.**

Myocardial ischemic events have been uncommonly reported in patients receiving irinotecan. In some cases, a causal association with administration of IRINOTECAN HYDROCHLORIDE INJECTION could not be excluded. See **Post-Market Adverse Drug Reactions.**

Endocrine and Metabolism

Hyponatremia has been reported in patients receiving irinotecan. This has usually been observed in patients with a history of dehydration or evidence of glucose intolerance prior to administration of IRINOTECAN HYDROCHLORIDE INJECTION (see also **Drug Interactions — Pharmacokinetics**).

Gastrointestinal

Diarrhea:

IRINOTECAN HYDROCHLORIDE INJECTION can induce both an early (occurring during or shortly after infusion of IRINOTECAN HYDROCHLORIDE INJECTION) and a late (generally occurring more than 24 hours after IRINOTECAN HYDROCHLORIDE INJECTION administration) form of diarrhea that appear to be mediated by different mechanisms.

Early onset diarrhea is cholinergic in nature. It is usually transient and only infrequently is severe. It may be accompanied by rhinitis, increased salivation, lacrimation, miosis, sweating, abdominal cramping, flushing and intestinal hypermotilities that can accelerate normal gastrointestinal charging. Early diarrhea can be alleviated by the use of atropine. Prophylactic or therapeutic administration of 0.25 to 1.0 mg of intravenous or subcutaneous atropine should be given with irinotecan (see DOSAGE AND ADMINISTRATION).

Late onset diarrhea can be prolonged, may lead to dehydration, electrolyte imbalance, or infection, and can be life-threatening. The mechanism of action of late onset diarrhea is unknown. Risk factors and late onset diarrhea occurred in 80% of patients and **late diarrhea should be treated promptly with loperamide.** Patients with diarrhea should be instructed to begin taking an over-the-counter loperamide at the first loose bowel movement. Loperamide should be given at initial dose of 4 mg followed by 2 mg every 4 hours. Loperamide should be continued for at least 12 hours after the last episode of diarrhea and should not be taken prophylactically. Loperamide should not be given for longer than 48 consecutive hours at these doses, because of the risk of paralytic ileus.

Inflammatory Bowel Disease and/or Bowel Obstruction:

Cases of colitis complicated by ulceration, bleeding, sepsis, and infection have been observed. Cases of ileus without preceding colitis have also been reported. Patients experiencing ileus should receive prompt antibiotics support (see DRUG INTERACTIONS) and must not be treated with irinotecan until resolution of the bowel obstruction.

Hematology

IRINOTECAN HYDROCHLORIDE INJECTION is emetogenic (see ADVERSE REACTIONS). Premedication with anti-emetic agents should be considered when administering IRINOTECAN HYDROCHLORIDE INJECTION. In clinical studies with the weekly dosage schedule, this premedication has been shown to reduce the incidence of severe nausea and vomiting and the severity of emetic regard. Anti-emetic agents should be given on the day of treatment, starting at least 30 minutes before administration of IRINOTECAN HYDROCHLORIDE INJECTION. Physicians should also consider providing patients with anti-emetic agents for subsequent use as needed.

Hematologic

Irinotecan commonly causes neutropenia, leukopenia, and anemia, any of which may be severe and therefore should not be used in patients with severe bone marrow failure. Therapy with irinotecan should not be initiated in patients with a neutrophil count of less than 1,500 cells/mm³ or platelets below 13 x 10⁹/L. After the patient recovers to a neutrophil count of 1,500 cells/mm³ or greater, subsequent doses of irinotecan should be reduced depending upon the level of neutropenia observed (see DOSAGE AND ADMINISTRATION). Severe neutropenia resulting in deaths due to sepsis have been reported in patients treated with irinotecan. Neutropenic complications should be managed promptly (see DOSAGE AND ADMINISTRATION). Patients with UGT1A1 genotype associated with decreased UGT1A1 activity (see CLINICAL PHARMACOLOGY) are at increased risk of developing severe neutropenia.

In a study an increased risk of neutropenia was observed in patients homozygous for the UGT1A1*28 allele, which received single-agent IRINOTECAN HYDROCHLORIDE INJECTION at 350 mg/m² (see CLINICAL PHARMACOLOGY). Individuals with certain genotypes of the UGT1A1 gene (e.g. UGT1A1 *28/*28 genotype) have elevated UGT1A1 activity, which in turn results in lower levels of SN-38, the active metabolite of irinotecan (see ACTION AND CLINICAL PHARMACOLOGY – Pharmacokinetics).

Patients with the UGT1A1*28 allele, treated with combination regimens that deliver doses of irinotecan in the range of 150-180 mg/m², in combination with 5-FU/LV, the risk of grade 4 neutropenia was lower than in studies where irinotecan was administered at doses of 300-350 mg/m² as a single agent (see DOSAGE AND ADMINISTRATION).

As in the case with all patients, dose modifications should be considered based on individual patient tolerance to treatment (See DOSAGE AND ADMINISTRATION).

Hepatic/Biliary/Pancreatic

Increases in serum levels of liver enzymes and bilirubin have been reported in clinical trials and in the post-market period (see ADVERSE DRUG REACTIONS).

The use of irinotecan in patients with significant hepatic dysfunction has not been established. Irinotecan was not administered to patients with serum bilirubin > 35 µmol/L or transaminases > 3 times the upper limit of normal if no liver metastases, or transaminases > 5 times the upper limit of normal if liver metastases were present (see DOSAGE AND ADMINISTRATION).

In clinical trials of weekly dosage schedule, patients with moderately elevated serum total bilirubin levels (17-35 µmol/L) had a significantly greater likelihood of experiencing first-cycle grade 3 or 4 hematologic toxicities including neutropenia, than patients with serum total bilirubin less than 17 µmol/L. Patients with deficient glucuronidation of bilirubin, such as those with Gilbert's syndrome, may be at greater risk of developing hematologic toxicity (see DOSAGE AND ADMINISTRATION). An association between baseline bilirubin elevations and an increased risk of severe toxicity has been observed in studies of the weekly dosage schedule (see DOSAGE AND ADMINISTRATION).

Immune

Hypersensitivity reactions including severe anaphylactic or anaphylactoid reactions have been reported (see ADVERSE REACTIONS).

Administration of live or live attenuated vaccines in patients immunocompromised by chemotherapeutic agents including irinotecan, may result in serious or fatal infections. Vaccination with a live vaccine should be avoided in patients receiving irinotecan. Killed or inactivated vaccines may be administered; however, the response to such vaccines may be diminished.

Neurologic

Speech disorders

There have been post-market reports of speech disorders (e.g. dysarthria, stuttering, change in speech quality, loss of speech) in patients receiving irinotecan. In some cases, including those in clinical trials, the speech disorder occurred in the context of other symptoms such as numbness or tingling of the tongue or mouth or symptoms attributed to cholinergic syndrome, hypersensitivity/allergy, cerebrovascular events, or intracranial neoplasia. In some cases, however, the speech disorder occurred in the absence of other symptoms and recurred with subsequent infusions of irinotecan. In some cases, the speech disorder resolved within minutes to hours after the infusion was stopped and resolved spontaneously within minutes to hours following discontinuation of the irinotecan infusion. The cause of speech disorders in patients receiving irinotecan has not been determined.

Driving and operating machinery

Patients should be warned about the potential for fatigue, dizziness or visual disturbances which may occur following the administration of irinotecan, and advised not to drive or operate machinery if these symptoms occur.

Renal

Increases in serum creatinine or blood urea nitrogen have been observed. Rare cases of renal insufficiency or failure have been reported in patients receiving irinotecan. Renal failure may be due to complications of infection or to dehydration related to nausea, vomiting and/or diarrhea, which are common and sometimes severe adverse events following irinotecan treatment. Rare instances of renal dysfunction due to tumour lysis syndrome have also been reported.

The influence of renal insufficiency on the pharmacokinetics of irinotecan has not been evaluated.

Respiratory

Interstitial pulmonary disease presenting as pulmonary infiltrates is uncommon during irinotecan therapy (see ADVERSE REACTIONS). Interstitial pulmonary disease can be fatal. Risk factors possibly associated with the development of interstitial pulmonary disease include pre-existing lung disease, concomitant chest radiation therapy, radiation therapy and/or chemotherapy before. Patients with risk factors should be closely monitored for respiratory symptoms before and during irinotecan therapy.

Special Populations

Pregnant Women:

Irinotecan has been shown to be embryotoxic in rats and rabbits at a dose of 6 mg/kg/day. It is teratogenic in rats at doses greater than 12 mg/kg/day, and in rabbits at 6 mg/kg/day. Treatment-related changes in the fetuses included external and visceral abnormalities, skeletal variations and abnormalities. Irinotecan may cause fetal harm when administered to a pregnant woman. If the drug is used during pregnancy, or if the patient becomes pregnant while receiving this drug, the patient should be apprised of the potential hazard to the fetus. Women of childbearing potential should be advised to avoid becoming pregnant while receiving IRINOTECAN HYDROCHLORIDE INJECTION.

Nursing Women:

In rats, radioactivity appeared in the milk within 5 minutes of intravenous administration of radiolabeled irinotecan and was concentrated up to 65-fold at 4 hours after administration relative to plasma concentrations (see DETAILED PHARMACOLOGY – Pharmacokinetics). It is not known whether irinotecan is excreted in human milk. Because many drugs are excreted in human milk and because of the potential for serious adverse reactions in nursing infants, it is recommended that nursing be discontinued when receiving therapy with IRINOTECAN HYDROCHLORIDE INJECTION.

Pediatrics:

The safety and effectiveness of irinotecan in the pediatric population have not been established.

Geriatrics:

Patients greater than 65 years of age should be closely monitored because of a greater risk of late severe toxicity (see DOSAGE AND ADMINISTRATION, CLINICAL TRIALS). The safety and efficacy of IRINOTECAN HYDROCHLORIDE INJECTION have not been established for the once-every-3-week-dosage schedule should be 300 mg/m² (see DOSAGE AND ADMINISTRATION).

Monitoring and Laboratory Tests

Careful monitoring of white blood cell count with differential, hemoglobin and platelet count is recommended before each dose of IRINOTECAN HYDROCHLORIDE INJECTION. Patients receiving irinotecan should be monitored before initiation of treatment and monthly, or as clinically indicated (See ADVERSE DRUG REACTIONS).

ADVERSE REACTIONS

Adverse Drug Reaction Overview

Gastrointestinal

Nausea, vomiting and diarrhea are common adverse events following treatment with IRINOTECAN HYDROCHLORIDE INJECTION and can be severe. When observed, nausea and vomiting usually occurred during the first 24 hours following administration. In clinical trials of the weekly dosage schedule, the median time to the onset of late diarrhea was 5 days after irinotecan infusion. In the clinical studies evaluating the weekly dosage schedule, the median time to onset of late diarrhea was 11 days following administration of irinotecan. All grade late diarrhea occurred in approximately 80% in this patient population. For patients on the 125 mg/m² weekly dose, the median duration of any grade late diarrhea was 4 days. The median duration was over 7 days for those patients reporting grade 3 or 4 diarrhea who were hospitalized at least once for diarrhea.

In a retrospective analysis, it has been shown that the frequency of grade 3 and 4 late diarrhea by age was significantly greater in patients ≥ 65 years than in patients < 65 years of age. However, results from a prospective study treating patients with metastatic colorectal cancer refractory to one 5-FU-based chemotherapeutic regimen, with a 125 mg/m² weekly dosage schedule of irinotecan (with 2 weeks off) did not demonstrate any statistically significant difference in the rate of treatment-related grade 3 or 4 diarrhea in patients ≥ 65 years of age compared to patients < 65 years of age. It should be noted however, that a statistically significant increase in the incidence of treatment-emergent early diarrhea in patients ≥ 65 years of age versus patients < 65 years of age was found. Furthermore, a 10% reduction in median relative weekly dose intensity was required in patients ≥ 65 years versus patients < 65 years in order to achieve tolerability. In the same prospective study, patients ≥ 65 years of age were found to have a significantly greater risk of treatment discontinuation due to adverse events, as well as a statistically significant increase in the incidence of treatment-emergent early diarrhea in patients ≥ 65 years of age versus patients < 65 years of age was found. Furthermore, a 10% reduction in median relative weekly dose intensity was required in patients ≥ 65 years versus patients < 65 years in order to achieve tolerability. In the same prospective study, patients ≥ 65 years of age were found to have a significantly greater risk of treatment discontinuation due to adverse events, as well as a statistically significant increase in the incidence of treatment-emergent early diarrhea in patients ≥ 65 years of age versus patients < 65 years of age was found.

Hematology

Typical adverse hematologic events of irinotecan included neutropenia, leukopenia (including lymphocytopenia), and anemia. Serious thrombocytopenia is uncommon. When evaluated in the trials of weekly administration, the frequency of grade 3 or 4 neutropenia was significantly increased in patients who had prior pelvic or abdominal irradiation. In the clinical studies evaluating the weekly dosage schedule, the median time to nadir was 15 days following administration. The nadir (grade 4 greater) occurred in 3.0% of the patients. Prior drug therapy did not appear to affect the incidence of neutropenia. NCI grade 3 or 4 anemia was noted in 6.9% of the patients. Blood transfusions were given to 9.9% of the patients. There was no significant differences in the frequency of grade 3 and 4 neutropenia by age or gender (see WARNINGS AND PRECAUTIONS, DOSAGE AND ADMINISTRATION). In a retrospective analysis of patients treated with the 125 mg/m² weekly dosage schedule, there is some information that patients with poor performance status may be at an increased risk of neutropenia or diarrhea.

Individuals with certain polymorphs of UGT1A1, such as UGT1A1*28, may have reduced activity of this enzyme. Approximately 10% of the North American population is homozygous for the UGT1A1*28 allele (also referred to as UGT1A1 7/7 genotype).

In a single arm study of 66 patients with solid tumors or lymphomas who received irinotecan as a single agent at 350 mg/m² every 3 weeks, a higher incidence of grade 4 neutropenia in patients with UGT1A1 *28/*28 genotype had grade 4 neutropenia versus 3 out of 24 patients with the UGT1A1 *1/*28 genotype, and 0 out of 29 patients with the UGT1A1 *1/*1 genotype (See WARNINGS AND PRECAUTIONS).

In a study which compared irinotecan (100-125 mg/m²) in combination with bolus 5-FU/LV (IFL regimen), oxaliplatin (85 mg/m²) in combination with irinotecan/5-FU/LV (FOLFOX6 regimen) and a combination of oxaliplatin (85 mg/m²) plus irinotecan (200 mg/m²) (IROX regimen), the incidence of grade 4 neutropenia is shown in the table below.

Rates of Grade 4 Neutropenia*† as Genotype by Treatment

UGT1A1 Genotype	IFL‡ n=109	FOLFOX6§ n=205	IROX¶ n=103
6/6	6.8% (3/44)	19.4% (36/134)	9.8% (5/52)
6/7	11.1% (6/54)	22.2% (28/126)	15.0% (8/40)
7/7	16.2% (2/11)	36.0% (9/25)	54.5% (6/11)

* IFL, Irinotecan 100-125 mg/m² followed by bolus 5-FU (500 mg/m²) and 5-FU 450 mg/m² given as a bolus on days 1, 8, 15, and 22 followed by a 2-week rest in repeated 6-week cycles.

† FOLFOX6, oxaliplatin 85 mg/m² in combination with irinotecan on days 1 and 2, and 5-FU 400 mg/m² given as a bolus followed by 600 mg/m² given as a 22 hr continuous infusion on days 1 and 2 in combination in repeated 2-week cycles.

‡ oxaliplatin 85 mg/m² followed by irinotecan 200 mg/m² administered on day 1 of repeated 3-week cycles.

‖ Fisher's exact test, IFL vs. FOLFOX6, p=0.0001; IFL vs. IROX, p=0.0001.

Pharmacogenetic studies have also suggested that the pharmacokinetic assessment of toxicity in patients with UGT1A1*28 genotype and reduced UGT1A1 activity, may be altered. Patients homozygous with the UGT1A1 *28/*28 genotype had grade 4 neutropenia, versus 6 out of 11 patients with the UGT1A1 *1/*28 and 2 out of 14 for patients with the UGT1A1 *1/*1 genotype.

Whole Body

Asthenia, fever, and abdominal pain are generally the most common events of this type.

Cholinergic Symptoms

Patients may have cholinergic symptoms of rhinitis, increased salivation, miosis, lacrimation, diaphoresis, flushing, and intestinal hypermotilities that can cause abdominal cramping and early diarrhea. If these symptoms occur, they maybe prevented during or shortly after drug infusion. They are thought to be related to the anticholinesterase activity of the irinotecan parent compound and are usually of short duration and rarely severe. Patients with asthma may be more susceptible to development of the cholinergic peak irinotecan serum levels than to parental administration.

Hepatic

In the clinical studies evaluating the weekly dosage schedule, NCI grade 3 or 4 liver enzyme abnormalities were observed in less than 10% of patients. These events typically occur in patients with known hepatic metastases.

Dermatologic

Alopecia has been reported during treatment with irinotecan. Rashes have also been reported but did not result in discontinuation of treatment.

Respiratory

Severe pulmonary events are infrequent. Early effects such as dyspnea have been reported (see WARNINGS AND PRECAUTIONS). In the clinical studies evaluating the weekly dosage schedule, over half the patients with dyspnea had lung metastases. The extent to which malignant pulmonary obstruction or other pre-existing lung disease may have contributed to dyspnea in these patients is unknown.

Potentially life threatening interstitial disease presenting with dyspnea, fever and pulmonary infiltrates (a noninfectious pattern on chest x-ray) is uncommon during irinotecan therapy. Usually seen in Japanese studies the combination of irinotecan to these events was difficult to assess because these patients also had lung tumours and some had pre-existing non-malignant pulmonary disease.

Neurologic

Insomnia and dizziness can occur, but are not usually considered to be directly related to the administration of irinotecan. Dizziness may sometimes represent symptomatic evidence of orthostatic hypotension in patients with dehydration (see WARNINGS AND PRECAUTIONS).

Cardiovascular

Vasodilation (flushing) may occur during administration of irinotecan. Bradycardia may also occur, but has not required intervention. These effects have been attributed to the cholinergic syndrome sometimes observed during or shortly after infusion of irinotecan. All thromboembolic events, when considered together, have commonly been observed in patients receiving irinotecan (see WARNINGS AND PRECAUTIONS, Cardiovascular). The specific cause of these events has not been determined.

Hypersensitivity

Hypersensitivity reactions including severe anaphylactic or anaphylactoid reactions have been observed (see WARNINGS AND PRECAUTIONS).

Clinical Trial Adverse Drug Reactions

Combination-Agent (Irinotecan/5-FU/LV) Therapy:

A total of 955 patients with metastatic colorectal cancer received regimens of irinotecan in combination with 5-FU/LV. 5-FU/LV alone, or irinotecan alone. In the two phase 3 studies, 370 patients received irinotecan in combination with 5-FU/LV, 362 patients received 5-FU/LV alone, and 223 patients received irinotecan alone (see Table 5 in DOSAGE AND ADMINISTRATION for dosage and schedule information).

In Study 1, 49 (13%) patients died within 30 days of last study treatment. 21 (9.3%) received irinotecan in combination with 5-FU/LV, 15 (8.4%) received 5-FU/LV alone, and 13 (5.8%) received irinotecan alone. Deaths potentially related to treatment occurred in 2 (0.9%) patients who received irinotecan in combination with 5-FU/LV (2 neutropenic fever/sepsis), 3 (1.4%) patients who received 5-FU/LV alone (1 neutropenic fever/sepsis, 1 CNS bleeding during thrombocytopenia, 1 unknown) and 1 (0.4%) patient who received irinotecan alone (neutropenic fever/sepsis). Deaths that occurred within 60 days of first study treatment were reported for 15 (6.7%) patients who received irinotecan in combination with 5FU/LV, 16 (13%) patients who received 5FU/LV alone and 15 (6.7%) patients who received irinotecan alone. Treatment-related deaths within 60 days were reported for 7 (3.0%) patients who received irinotecan in combination with 5-FU/LV, 16 (6.4%) patients who received 5-FU/LV alone and 1 (0.4%) patient who received irinotecan alone.

In Study 1, data on hospitalizations for any reason (hospitalization required as a consequence of chemotherapy-induced adverse events and hospitalizations that may have resulted from complications of the disease) were collected for all patients. Overall, 144 (62%) patients in Study 1 of 225 who received irinotecan in combination with 5-FU/LV were hospitalized. Sixty-eight (30.2%) patients who received 5-FU/LV alone and 54 (24.2%) patients who received irinotecan alone were hospitalized more than once. Forty-one (18.4%) patients treated with irinotecan/5-FU/LV were hospitalized. Twenty-one (18%) patients were hospitalized once, 21 (9.4%) were hospitalized twice and 7 (3.1%) were hospitalized 3 times. Forty-four (19.7%) patients and 27 (12%) patients with 5-FU/LV were hospitalized. Sixty (27.4%) patients were hospitalized once, 29 (13%) patients were hospitalized twice and 15 (6.7%) were hospitalized 3 times.

In Study 2, 10 (3.5%) patients died within 30 days of last study treatment: 6 (4.1%) received irinotecan alone, 3 (2.1%) received 5-FU/LV alone, and 1 (0.7%) received both irinotecan and 5-FU/LV. Deaths potentially related to treatment occurred in 4 patients: 2 (1.4%) received irinotecan alone (1 neutropenic fever/sepsis, 1 unknown), 1 (0.7%) received 5-FU/LV alone (1 neutropenic fever/sepsis), and 1 (0.7%) received both irinotecan and 5-FU/LV (1 diarrhea, neutropenic fever/sepsis). Deaths that occurred within 60 days of first study treatment were reported for 15 (10.4%) patients who received irinotecan alone, 7 (4.9%) patients who received 5-FU/LV alone, and 15 (10.4%) patients who received both irinotecan and 5-FU/LV. Treatment-related deaths within 60 days were reported for 4 (2.8%) patients who received irinotecan alone, 1 (0.7%) patient who received 5-FU/LV alone, and 3 (2.1%) patients who received both irinotecan and 5-FU/LV.

In Study 2, data on hospitalizations for any reason (hospitalization required as a consequence of chemotherapy-induced adverse events and hospitalizations that may have resulted from complications of the disease) were collected for all patients. Overall, 103 (71%) patients in Study 2 of 145 who received irinotecan in combination with 5-FU/LV were hospitalized. Sixty-eight (30.2%) patients who received 5-FU/LV alone and 54 (24.2%) patients who received irinotecan alone were hospitalized more than once. Forty-one (28.3%) patients treated with irinotecan/5-FU/LV were hospitalized. Twenty-one (14%) patients were hospitalized once, 21 (9.4%) were hospitalized twice and 7 (3.1%) were hospitalized 3 times. Forty-four (19.7%) patients and 27 (12%) patients with 5-FU/LV were hospitalized. Sixty (27.4%) patients were hospitalized once, 29 (13%) patients were hospitalized twice and 15 (6.7%) were hospitalized 3 times.

In Study 2, 10 (3.5%) patients died within 30 days of last study treatment: 6 (4.1%) received irinotecan alone, 3 (2.1%) received 5-FU/LV alone, and 1 (0.7%) received both irinotecan and 5-FU/LV. Deaths potentially related to treatment occurred in 4 patients: 2 (1.4%) received irinotecan alone (1 neutropenic fever/sepsis, 1 unknown), 1 (0.7%) received 5-FU/LV alone (1 neutropenic fever/sepsis), and 1 (0.7%) received both irinotecan and 5-FU/LV (1 diarrhea, neutropenic fever/sepsis). Deaths that occurred within 60 days of first study treatment were reported for 15 (10.4%) patients who received irinotecan alone, 7 (4.9%) patients who received 5-FU/LV alone, and 15 (10.4%) patients who received both irinotecan and 5-FU/LV. Treatment-related deaths within 60 days were reported for 4 (2.8%) patients who received irinotecan alone, 1 (0.7%) patient who received 5-FU/LV alone, and 3 (2.1%) patients who received both irinotecan and 5-FU/LV.

Table 1. Study 1: Percent (%) of Patients Experiencing Clinically Relevant Adverse Events in Combination Therapies*

Adverse Event	Study 1					
	Irinotecan + Bolus 5-FU/LV q 4 weeks N=226		Bolus 5-FU/LV q 4 weeks N=219		Irinotecan weekly q 4 weeks N=223	
	Grade 1-4	Grade 3-4	Grade 1-4	Grade 3-4	Grade 1-4	Grade 3-4
TOTAL Adverse Events	100	53.3	100	45.7	99.6	45.7
GASTROINTESTINAL						
Diarrhea						
late	84.9	22.7	69.4	13.2	83.0	31.5
grade 3	–	10.1	–	5.0	–	16.4
grade 4	–	7.6	–	7.3	–	12.6
early	45.6	4.9	13.2	0.5	41.3	6.7
Nausea	79.1	15.6	67.6	8.2	81.6	16.1
Abdominal pain	63.1	10.2	50.2	4.1	61.4	11.2
Vomiting	60.4	9.7	48.1	4.1	62.8	12.1
Anorexia	34.2	2.7	42.0	1.8	43.5	4.5
Constipation	41.3	3.1	31.5	1.8	32.3	0.4
Mucositis	32.4	4.4	26.3	1.4	29.6	2.2
GASTROINTESTINAL						
Diarrhea						
late	96.9	53.8	98.6	66.7	96.4	31.4
grade 3	–	29.8	–	23.7	–	19.3
grade 4	–	24.0	–	42.5	–	12.1
Leukopenia	96.9	37.8	98.6	23.3	96.4	21.5
Anemia	96.9	8.4	98.6	6.8	96.4	4.5
Neutropenic fever	–	7.1	–	14.6	–	5.8
Thrombocytopenia	–	2.6	98.6	2.7	96.4	1.7
Neutropenic infection	–	1.8	–	–	–	2.2
BODY AS A WHOLE						
Asthenia	70.2	19.5	64.4	11.9	69.1	13.9
Pain	30.7	2.2	34.8	3.2	36.8	3.6
Fever	42.2	1.7	32.4	3.6	43.5	0.4
Infection	22.2	0	16.0	1.4	13.9	0.4
METABOLIC & NUTRITIONAL						
† Bilirubin	87.6	7.1	92.2	8.2	83.9	7.2
DERMATOLOGIC						
Exfoliative dermatitis	0.9	0	3.2	0.5	0	0
Rash	19.1	0	26.5	0.9	14.3	0.4
Alopecia‡	43.1	–	36.5	–	45.7	–
RESPIRATORY						
Dyspnea	27.6	6.3	16.0	0.5	22.0	2.2
Cough	26.7	1.3	18.3	0	20.2	0.4
Pneumonia	6.2	2.7	1.4	1.0	4.5	1.3
NEUROLOGIC						
Dizziness	23.1	1.3	16.4	0	21.1	1.8
Somnolence	12.4	1.8	4.6	1.8	9.4	1.3
Confusion	7.1	1.8	4.1	0	2.7	0
CARDIOVASCULAR						
Vasodilatation	9.3	0.9	5.0	0	9.0	0
Syncope	5.8	2.3	0.5	0.5	5.8	1.7
Thromboembolic phenomena	9.3	–	11.4	5.4	–	–

* Severity of adverse events based on NCI CTC criteria 1.0.

† Bilirubin > 1.5 x ULN

‡ Includes angina pectoris, arterial thrombosis, cerebral infarct, cerebrovascular accident, deep thrombophlebitis, embolus (over extremity), heart arrest, myocardial infarct, myocardial ischemia, peripheral vascular disorder, pulmonary embolus, sudden death, thrombophlebitis, venous thrombosis.

§ Includes cholinergic syndrome, dehydration, hypotension, hypovolemia, intestinal obstruction, pancreatitis and sudden death.

Table 2. Study 2: Percent (%) of Patients Experiencing Clinically Relevant Adverse Events in Combination Therapies*

Adverse Event	Study 2			
	Irinotecan +5-FU/LV initiated d1&2 q 2 weeks N=145		5-FU/LV infusional d1&2 q 2 weeks N=143	
	Grade 1-4	Grade 3-4	Grade 1-4	Grade 3-4
GASTROINTESTINAL				
Diarrhea				
late	72.4	14.4	44.8	6.3
grade 3	–	10.3	–	4.2
grade 4	–	4.1	–	2.1
Cholinergic syndrome†	28.3	1.4	0.7	0
Nausea	66.9	2.1	55.2	3.5
Abdominal pain	17.2	2.1	14.8	0.7
Vomiting	44.8	3.5	32.2	2.8
Anorexia	35.2	2.1	18.9	0.7
Constipation	30.3	0.7	25.2	1.4
Mucositis	40	4.1	32.2	2.8
HEMATOLOGIC				
Neutropenia	82.5	–	46.2	47.9	13.4	–
grade 3	–	–	36.4	–	12.7	8.9
grade 4	–	–	9.8	–	0.7	39.0
Leukopenia	81.3	–	17.4	–	42	3.5
Anemia	97.2	–	2.1	–	90.9	2.1
Neutropenic fever	–	–	3.4	–	0.7

^CHLORHYDRATE D'IRINOTÉCAN POUR INJECTION

20 mg/mL de chlorhydrate d'irinotécan (trihydraté)

PARTIE I : RENSEIGNEMENTS POUR LES PROFESSIONNELS DE LA SANTÉ

RENSEIGNEMENTS SOMMAIRES SUR LE PRODUIT

Voie d'administration	Présentation et teneur	Excipients d'importance clinique
Intraveineuse	Solution dosée à 20 mg/mL	Sorbitol, Voir la section PRÉSENTATION, COMPOSITION ET CONDITIONNEMENT pour connaître la liste complète des ingrédients.

INDICATIONS ET USAGE CLINIQUE

Le CHLORHYDRATE D'IRINOTÉCAN POUR INJECTION (*chlorhydrate d'irinotécan*) est indiqué en association avec d'autres agents pour le traitement des patients atteints d'un cancer colorectal du côté droit ou modéré.

Le CHLORHYDRATE D'IRINOTÉCAN POUR INJECTION est également indiqué en monothérapie en présence d'un cancer métastatique du côté droit ou modéré qui a évolué ou a évolué pendant ou après le 5-fluorouracile.

Le CHLORHYDRATE D'IRINOTÉCAN POUR INJECTION ne doit être administré que sous la surveillance d'un médecin connaissant bien l'emploi des agents de chimiothérapie anticancéreux. Il faut disposer d'installations appropriées et d'un personnel médical compétent d'une nature et d'une qualité adéquates pour traiter l'éventuel acide à un établissement de soins équipé pour faire face à ce genre de situation.

Personnes âgées : Des données issues d'études cliniques et de la pharmacocinétique semblent indiquer que les patients âgés de 65 ans ou plus doivent être suivis de près en raison d'un risque accru de diarrhée tardive dans cette population. En fonction du schéma posologique sélectionné, il se peut qu'il faille suivre des recommandations posologiques particulières pour ce groupe d'âge (voir POSOLOGIE ET MODE D'ADMINISTRATION – Posologie recommandée, Ajustements posologiques, ET MODE D'ADMINISTRATION et ESSAIS CLINIQUES).

Enfants : L'innocuité et l'efficacité du CHLORHYDRATE D'IRINOTÉCAN POUR INJECTION chez l'enfant n'ont pas été établies (voir la section MISES EN GARDE ET PRÉCAUTIONS).

CONTRE-INDICATIONS

Le CHLORHYDRATE D'IRINOTÉCAN POUR INJECTION est contre-indiqué chez les patients qui présentent une hypersensibilité à ce médicament ou à l'un des composants du produit ou du contenant.

Voir la rubrique PRÉSENTATION, COMPOSITION ET CONDITIONNEMENT pour connaître la liste complète des ingrédients.

L'administration concomitante du CHLORHYDRATE D'IRINOTÉCAN POUR INJECTION et d'inducteurs puissants de l'isoenzyme CYP3A4, tels que la phénytoïne, la carbamazépine, le phénobarbital ou la CYP3A4, est contre-indiquée puisque ces médicaments peuvent réduire la biodisponibilité et la métabolite actif SN-38 et ainsi augmenter la toxicité de l'irinotécan.

Chez les patients recevant de l'irinotécan et de la bilirubinémie en concomitance, l'exposition au SN-38 a augmenté d'environ 110 %. Les patients doivent cesser de prendre du kétoconazole au moins une semaine avant le début du traitement par l'irinotécan. Si l'on ne peut éviter l'administration concomitante médicamenteuses avec d'autres inducteurs de la CYP3A4, il faut envisager d'augmenter la dose (voir la rubrique MISES EN GARDE ET PRÉCAUTIONS).

L'administration du CHLORHYDRATE D'IRINOTÉCAN POUR INJECTION est contre-indiquée chez les patients présentant une infection hématologique ou fécale, car ce produit contient du sorbitol.

MISES EN GARDE ET PRÉCAUTIONS

Mises en garde et précautions importantes

- Le CHLORHYDRATE D'IRINOTÉCAN POUR INJECTION ne doit être administré que sous la supervision d'un médecin possédant une expérience en chimiothérapie.
- Le CHLORHYDRATE D'IRINOTÉCAN POUR INJECTION peut provoquer des diarrhées graves. Il faut donc surveiller étroitement les patients présentant ce symptôme (voir les rubriques MISES EN GARDE ET PRÉCAUTIONS, Gastro-intestinal, et Aperçu des effets indésirables du médicament).
- Le CHLORHYDRATE D'IRINOTÉCAN POUR INJECTION peut provoquer une syphilite, des colites ulcéreuses et ischémiques, un iléus et des perforations intestinales (voir les rubriques MISES EN GARDE ET PRÉCAUTIONS, Gastro-intestinal, et EFFETS INDÉSIRABLES). Effets indésirables du médicament, Aperçu des effets indésirables du médicament).

Généralités

Patients exposés à un risque particulier

Les médecins doivent surveiller avec une extrême vigilance les effets du CHLORHYDRATE D'IRINOTÉCAN POUR INJECTION chez les patients qui présentent un risque accru de toxicité. Le CHLORHYDRATE D'IRINOTÉCAN POUR INJECTION ne doit pas être administré à des patients ayant un indice fonctionnel de 3 ou de 4 (selon la cote de l'échelle de performance du groupe d'oncologie de l'Amérique du Nord) ou à des patients présentant une maladie hépatique importante (voir la rubrique MISES EN GARDE ET PRÉCAUTIONS, Hépatique).

Système endocrinien/métabolisme

Des cas d'hyperglycémie ont été rapportés chez des patients traités à l'irinotécan. Ces cas ont généralement été observés chez des patients présentant des facteurs de risque préexistants tels qu'un glucose avant l'administration du CHLORHYDRATE D'IRINOTÉCAN POUR INJECTION (voir INTERACTIONS MÉDICAMENTEUSES, Interactions médicament-médicament, Dexamethasone).

Appareil digestif

Diarrhée

Le CHLORHYDRATE D'IRINOTÉCAN POUR INJECTION peut provoquer une diarrhée précoce (survenant pendant ou peu après la perfusion de CHLORHYDRATE D'IRINOTÉCAN POUR INJECTION) ou tardive (survenant généralement plus de 24 heures après l'administration du CHLORHYDRATE D'IRINOTÉCAN POUR INJECTION). Il faut surveiller étroitement les patients présentant une diarrhée précoce et d'origine cholinergique. Elle est généralement passagère et n'est que rarement grave. Elle peut être accompagnée de rhinite, de larmoiement, de myosis, de bradycardie, de tachycardie, de transpiration, de salivation, de crampes abdominales et de symptômes gastro-intestinaux. Il faut donc administrer de l'atropine à titre préventif aux patients présentant ces symptômes. La diarrhée précoce peut être soulagée par l'administration d'atropine. L'administration préventive d'atropine doit être envisagée à une dose de 0,25 à 1,0 mg d'atropine, par voie intraveineuse ou sous-cutanée, dès que l'irinotécan doit être administré à des patients qui ont déjà présenté ce symptôme (voir la rubrique POSOLOGIE ET MODE D'ADMINISTRATION).

La diarrhée tardive peut être très grave. Elle peut entraîner une déshydratation, un déséquilibre électrolytique ou une infection, ou encore la mort si elle n'est pas traitée. Elle peut aussi avoir des répercussions sur l'état physiologique de la flore intestinale. Elle survient généralement plus tard, en général après plus de 24 heures après l'administration du CHLORHYDRATE D'IRINOTÉCAN POUR INJECTION (voir la rubrique MISES EN GARDE ET PRÉCAUTIONS, Gastro-intestinal). Les patients atteints de diarrhée doivent faire l'objet d'une surveillance étroite et recevoir rapidement un traitement hydro-électrolytique en cas de déshydratation. Les patients doivent suivre une antibiothérapie (voir la rubrique MISES EN GARDE ET PRÉCAUTIONS, Gastro-intestinal) en présence de fièvre ou de neutropénie grave. Après la première séance de traitement, la séance suivante ne doit pas commencer tant que la transit intestinal du patient n'est pas redevenu ce qu'il était avant le traitement sans prise d'un antidiarrhéique pendant au moins 24 heures, et tant que le patient ne présente pas de point de vue clinique (grade ≤ 2). Il faut réduire les doses d'irinotécan administrées par la suite (voir la rubrique POSOLOGIE ET MODE D'ADMINISTRATION).

Traitement de la diarrhée tardive : Au début de la diarrhée tardive, il faut encourager les patients à ingérer une quantité suffisante de hydratation et de bien lui en expliquer le mode d'emploi. Il est recommandé de prendre la loperamide par voie orale après le premier épisode de selles molles et de diarrhée et d'utiliser une dose supérieure à la dose usuelle. On ne conseille pas de prétraitement au loperamide avant le traitement par l'irinotécan. Il faut conseiller aux patients de commencer à prendre 4 mg de loperamide au premier signe de diarrhée, puis 2 mg toutes les 2 heures. Il faut poursuivre cette dose jusqu'à ce que la diarrhée ait disparu depuis au moins 12 heures. Pendant la nuit, le patient peut prendre 4 mg de loperamide toutes les 4 heures. Le traitement par la loperamide à la dose prescrite pendant plus de 48 heures consécutives est déconseillé à cause du risque d'iléus paralytique.

Maladie inflammatoire de l'intestin et/ou occlusion intestinale

On a observé des cas de colite compatible d'ulcération, de saignements, d'occlusion intestinale et d'infection. Des cas d'occlusion intestinale ont été rapportés chez des patients qui avaient reçu la patient atteint d'une occlusion intestinale doit rapidement recevoir des antibiotiques dans le cadre d'un traitement approprié et faire l'objet d'une surveillance étroite. Il faut arrêter le traitement par l'irinotécan chez les patients qui présentent une diarrhée sévère ou une occlusion intestinale.

Nausées et vomissements

Le CHLORHYDRATE D'IRINOTÉCAN POUR INJECTION a des effets émétiques (voir la rubrique EFFETS INDÉSIRABLES). L'administration d'un antiémétique est recommandée avant le début du traitement au CHLORHYDRATE D'IRINOTÉCAN POUR INJECTION. Il faut prescrire aux patients un traitement contre la sphère d'administration hebdomadaire, cette démarche comportait le plus souvent de la dexaméthasone à 8 mg, par voie intraveineuse ou orale, 30 minutes à 1 heure avant l'administration du CHLORHYDRATE D'IRINOTÉCAN POUR INJECTION. Les patients ayant présenté des vomissements tardifs ou persistants après un premier traitement doivent également recevoir de la dexaméthasone (par exemple, de 4 mg à 8 mg, deux fois par jour) pendant 2 à 4 jours. Les patients ayant présenté des nausées ou des vomissements malgré l'administration du CHLORHYDRATE D'IRINOTÉCAN POUR INJECTION. Le médecin doit également considérer la prescrire un traitement antiémétique que le patient pourra prendre au besoin par la suite.

Appareil circulatoire

L'irinotécan est une cause fréquente de neutropénie, de leucopénie et d'anémie, des affections toutes susceptibles de se compliquer d'une infection ou d'une hémorragie. La neutropénie est le plus souvent modérée et réversible. Elle est rarement grave. Le traitement à l'irinotécan ne doit pas être entrepris chez les patients dont le taux de neutrophile fibrillo ou si le nombre de polynucléaires neutrophiles chute au-dessous de 1,5 x 10⁹/L. Lorsque le nombre de polynucléaires neutrophiles chute au-dessous de 1 pour 1,5 x 10⁹/L, il faut réduire les doses subséquentes d'irinotécan suivant le degré de neutropénie observé (voir la rubrique POSOLOGIE ET MODE D'ADMINISTRATION). Il faut surveiller étroitement les patients présentant une neutropénie grave (moins de 0,5 x 10⁹/L pendant plus de 7 jours). Il faut également surveiller les patients présentant une neutropénie fébrile (définie par un nombre de polynucléaires neutrophiles absolu inférieur à 1,0 x 10⁹/L et une température supérieure à 38,5 °C) ou une infection documentée (voir la rubrique INTERACTIONS MÉDICAMENTEUSES). Lorsque le médecin doit considérer l'utilisation de tels facteurs en présence de neutropénie notable ou en cas de neutropénie prolongée (plus de 7 jours).

Au cours d'une étude, un risque accru de mortalité a été observé chez des patients hémorragiques (voir la rubrique MISES EN GARDE ET PRÉCAUTIONS, Appareil circulatoire). Dans cette étude, le CHLORHYDRATE D'IRINOTÉCAN POUR INJECTION, à une dose de 350 mg/m² (dose trop élevée), a été associé à une incidence accrue de décès par hémorragie gastro-intestinale et de décès par infection. Ces décès ont généralement été observés pendant les deux premières semaines de cycle, et il a été difficile d'évaluer dans quelle mesure l'irinotécan y a contribué. Les patients doivent également prendre connaissance de tumeurs pulmonaires et certains avaient une pneumopathie non cancéreuse préexistante.

Troubles du système nerveux

Des cas d'insomnie et d'étourdissements peuvent se produire, mais on ne connaît généralement pas leur lien de causalité avec l'irinotécan. Les patients doivent faire preuve de prudence jusqu'à ce que les étourdissements ou symptômes apparentés à l'hypertension orthostatique chez les patients en état de déshydratation à la suite de vomissements ou de diarrhée.

Troubles oculaires

Une vasodilatation (troubles vasomoteurs) peut se produire pendant l'administration de l'irinotécan. Les cas de bradycardie peuvent être traités par l'atropine. Il faut également prendre des mesures d'intervention au cours des études cliniques. Des effets indésirables oint été attribués au syndrome cholinergique qui survient pendant ou peu après l'administration d'irinotécan. Ces effets sont généralement de brève durée et peuvent être traités par l'atropine. Les patients doivent également être informés des troubles oculaires, tels que la diplopie, la vision trouble, la photophobie et la douleur oculaire, qui peuvent également survenir pendant ou après l'administration d'irinotécan.

Hypersensibilité

On a signalé des réactions d'hypersensibilité, dont des réactions anaphylactiques ou anaphylactoïdes graves (voir la rubrique MISES EN GARDE ET PRÉCAUTIONS).

Effets indésirables du médicament observés au cours des essais cliniques

Traitement à l'association irinotécan/5-FU/LV

Chez les patients recevant un cancer métastatique du côlon ou du rectum qui ayant déjà schémas thérapeutiques comportant l'emploi de l'association irinotécan/5-FU/LV, du 5-FU/LV ou de l'irinotécan en monothérapie, l'innocuité de l'irinotécan a été évaluée dans le cadre d'une étude irinotécan/5-FU/LV 382 ont reçu le 5-FU/LV et 223 ont reçu l'irinotécan en monothérapie. Les patients ont reçu l'irinotécan à la dose de 350 mg/m² toutes les trois semaines ou le 5-FU/LV selon le schéma posologique hebdomadaire suivant : 500 mg/m² par voie i.v. chaque semaine pendant 6 semaines, suivi de 2 semaines de repos avant le cycle suivant. Le schéma posologique du 5-FU/LV était le suivant : 500 mg/m² par voie i.v. chaque semaine pendant 6 semaines, suivi de 2 semaines de repos avant le cycle suivant. Les caractéristiques démographiques étaient les suivantes :

Dans le bras irinotécan/5-FU/LV, 56 % des patients étaient des hommes et 44 % des femmes. Les études du système nerveux comprenaient les symptômes cholinergiques, l'asthénie et la dysfonctionnement de la vésicule biliaire.

Parmi les autres manifestations, citons les traumatismes accidentels, l'hypotonie/ligne, les syncopes, les vertiges et la perte de poids.

Effets indésirables du médicament signalés après la commercialisation du produit

Les effets suivants ont été signalés pendant l'utilisation thérapeutique de ce produit chez des patients après sa commercialisation.

Il se rares cas de colite, y compris de typhlite, de colite ulcéreuse et de colite ischémique, ont été signalés. Les complications graves comprenaient la perforation intestinale, l'hémorragie intestinale, la coma ou migration toxique, des ulcères, des saignements, une obstruction et une infection. De rares cas de perforation de l'intestin grêle ont également été signalés. L'occlusion intestinale s'est produite d'une cotte se sont aussi produits. Une antidiarrhéique doit être amorcée dans les plus brefs délais dès l'apparition de symptômes de diarrhée (voir la rubrique MISES EN GARDE ET PRÉCAUTIONS). Des cas de septicémie, parfois mortelle, ont été signalés.

Des hyponatrémie liée surtout à la diarrhée et aux vomissements a été observée dans de rares cas.

Des cas isolés de pneumopathie d'étiologie inconnue ont été signalés. Dans certains cas, des cas d'infiltrats pulmonaires, d'épanchement pleural, d'insuffisance respiratoire ou de détresse respiratoire aiguë ont été signalés. Des cas de pneumopathie interstitielle, parfois mortelle, ont été signalés.

De rares cas de syndrome de leucoencéphalopathie postérieure réversible (SLPR) ont été signalés. Les symptômes comprenaient des céphalées, des convulsions, des troubles de la conscience et des troubles visuels. Des examens complémentaires, tels que l'imagerie par résonance magnétique (IRM), ont été nécessaires pour confirmer le diagnostic. Les patients doivent faire l'objet d'une surveillance étroite et recevoir un traitement approprié.

Des cas d'anaphylaxie ont été signalés.

Des cas d'insuffisance rénale, d'hypertension ou de collapsus cardiovasculaire sont survenus chez des patients ayant présenté une déshydratation liée à la diarrhée et/ou à des vomissements, ou ayant souffert de septicémie (voir la rubrique MISES EN GARDE ET PRÉCAUTIONS).

Des effets précoces, tels que des contractions ou des crampes musculaires et une paresthésie, ont été signalés.

Les effets respiratoires graves sont rares. Les cas de pneumopathie interstitielle se manifestant par un infiltrat pulmonaire, un épanchement pleural, une insuffisance respiratoire ou un syndrome de détresse respiratoire aiguë ont été signalés (voir la rubrique MISES EN GARDE ET PRÉCAUTIONS).

On a observé des accidents ischémiques du myocarde, montés dans certains cas, chez des patients traités à l'irinotécan, y compris des infarctus du myocarde. Ces événements ont généralement survenu chez des patients présentant des facteurs de risque de maladie cardiaque et/ou étaient traités en concomitance par un autre antinéoplasique cytotoxique (voir la rubrique MISES EN GARDE ET PRÉCAUTIONS).

Des cas de syndrome de lyse tumorale, d'hyperkaliémie et d'hypocalcémie ont été signalés.

Des effets précoces, tels que des contractions ou des crampes musculaires et une paresthésie, ont été signalés.

INTERACTIONS MÉDICAMENTEUSES

Aperçu

L'irinotécan est transformé en un métabolite actif, le SN-38, par une carboxylestérase et il est ensuite éliminé par glucuronidation par l'UGT1A1. Le SN-38 et le SN-38 G sont éliminés par la bile et l'urine. Les études *in vitro* ont démontré que l'irinotécan est un substrat (voir la rubrique MODE D'ACTION ET PHARMACOLOGIE CLINIQUE, Pharmacocinétique). Des interactions pharmacocinétiques avec l'irinotécan, les métabolites actifs ou inactifs ont été observées chez l'humain et/ou l'animal. La plupart de ces interactions ont été attribuées à l'induction ou à l'inhibition de la CYP3A4, ou à l'inhibition de l'UGT1A1. Les études *in vitro* ont également démontré que l'irinotécan et ses métabolites sont des substrats de la glycoprotéine P (voir la rubrique MODE D'ACTION ET PHARMACOLOGIE CLINIQUE, Pharmacocinétique).

D'après les résultats d'études sur les interactions médicamenteuses *in vitro*, la biotransformation de l'irinotécan en son métabolite actif SN-38 par les carboxylestérases n'est pas inhibée par le fluorouracile (5-FU), les sels de platine ou la dexaméthasone. La phase d'essai sur l'administration de l'irinotécan, du 5-FU et de la leucovorine (LV) à 26 patients porteurs de tumeurs solides avancées, on a observé une augmentation de l'exposition au SN-38 et au SN-38 G, ainsi qu'une augmentation de la toxicité hématologique. Chez ces patients, l'administration concomitante de l'irinotécan et de l'irinotécan n'a pas modifié la pharmacocinétique du 5-FU. Les interactions médicamenteuses *in vivo* qu'*in vitro* n'ont été effectuée pour évaluer l'influence de l'irinotécan sur le métabolisme et le 5-FU.

L'administration concomitante de l'irinotécan avec des inhibiteurs de la CYP3A4 (tels que kétoconazole, érythromycine, ritonavir, indinavir, itraconazole, clarithromycine, fluconazole, amprénavir, nelfinavir et grapefruit) peut entraîner une augmentation de l'exposition au SN-38. Le pamplemousse et les biopsies des canaux calciques qui inhibent la CYP3A4 lorsqu'ils, plaidant, ou les médicaments qui inhibent la CYP3A4 doivent être évités. Une telle interaction a été observée chez des cancéreux recevant en concordance de l'irinotécan et du kétoconazole. Chez ces patients, l'ASC du SN-38 a été multipliée par 3,5 environ (IC 90 % : 2,81 à 4,66) alors que l'ASC du SN-38 G a été multipliée par 2,2 environ (IC 90 % : 1,81 à 2,79). Cette interaction a été observée chez des cancéreux recevant en concordance de l'irinotécan et du ritonavir, et de l'irinotécan et de la phénytoïne. Il faut aussi consulter les renseignements thérapeutiques sur les médicaments concomitants pour connaître les interactions médicamenteuses.

Interactions médicament-médicament

Interactions pharmacologiques

	Réf.*	Effet	Commentaire
Inhibiteurs de la CYP3A4			
Antifongiques azolés :			
Kétoconazole	EC	↑ environ 110 % du SN-38 ↑ environ 90 % du SNPC	Voir la rubrique CONTRE-INDICATIONS
Fluconazole, Itraconazole	T		
Cimétidine	T		
Antibiotiques de la classe des fluoroquinolones	T		
Ciprofloxacine, norfloxacine	T		
Antibiotiques de la classe des macrolides :	T		Voir la rubrique MISES EN GARDE ET PRÉCAUTIONS
Azithromycine, clarithromycine, érythromycine			
Bloqueurs des canaux calciques	T		
Diltiazem, vérapamil, nifédipine	T		
Jus de pamplemousse	T		
Sulfate d'atazanavir	T	Voir la monographie du médicament pour connaître les renseignements ci-après	
Inducteurs de la CYP3A4			**Risque d'efficacité moindre**
Anticonvulsivants			
Carbamazépine, phénobarbital, phénytoïne	ETC	↓ environ 65 % de l'irinotécan ↓ environ 75 % du SN-38 ↓ environ 40 % du SNPC	Voir la rubrique MISES EN GARDE ET PRÉCAUTIONS et Interactions médicament-herbe médicinale ci-dessous.
Glucocorticoïdes	C		
Rifampicine	T		

*Niveau de preuve. C = description de cas, EC = essai clinique, T = théorique

Le jous irinotécan approprié pour les patients prenant des médicaments ayant un effet, connu ou probable, sur la cinétique de l'irinotécan ou de ses métabolites. L'administration concomitante d'antifongiques azolés et d'irinotécan est contre-indiquée, et les patients doivent cesser de prendre du kétoconazole au moins une semaine avant le début du traitement par l'irinotécan. Si l'on ne peut éviter l'administration concomitante de médicaments qui sont des inducteurs puissants de la CYP3A4, tels que la phénytoïne, la carbamazépine ou le phénobarbital, il faut envisager d'augmenter la dose (voir la rubrique CONTRE-INDICATIONS). Les patients ne doivent pas faire de jus de pamplemousse durant le traitement. Si la prise d'anticonvulsants est nécessaire, le recours à ceux qui ne sont pas des inducteurs puissants de la CYP3A4 (comme le valproate de sodium) doit être envisagé au moins 1 semaine avant le début du traitement à l'irinotécan chez les patients qui ne prennent pas déjà d'anticonvulsants inducteurs. Il faut également éviter de prendre de la CYP3A4 et de l'UGT1A1, tels que le atazanavir ou le SN-38. Il est recommandé de consulter la monographie du médicament pour connaître les interactions médicamenteuses.

Interactions pharmacodynamiques

Antinéoplasiques : On peut s'attendre à ce que les effets indésirables du CHLORHYDRATE D'IRINOTÉCAN POUR INJECTION, tels que la dépression médullaire et la diarrhée, soient aggravés par l'administration concomitante d'autres antinéoplasiques (voir la rubrique MISES EN GARDE ET PRÉCAUTIONS).

Laxatifs : On peut s'attendre à ce que la fréquence ou l'intensité de la diarrhée soient aggravées par l'administration concomitante de laxatifs (voir la rubrique MISES EN GARDE ET PRÉCAUTIONS).

Diurétiques : L'administration concomitante de diurétiques peut entraîner une déshydratation due à des vomissements et/ou à la diarrhée entraînés par le CHLORHYDRATE D'IRINOTÉCAN POUR INJECTION. Il est possible que la probabilité de cet effet soit accrue par l'administration de diurétiques pendant l'administration du CHLORHYDRATE D'IRINOTÉCAN POUR INJECTION (voir la rubrique MISES EN GARDE ET PRÉCAUTIONS).

Dexaméthasone : On a observé une augmentation de l'hyperglycémie chez les patients traités à l'irinotécan. Cet effet a généralement été observé chez des patients présentant des facteurs de risque préexistants d'hyperglycémie, et certains patients ont présenté une hyperglycémie grave, et aucune complication n'a été signalée. Il faut donc surveiller étroitement le glucose avant l'administration du CHLORHYDRATE D'IRINOTÉCAN POUR INJECTION chez les patients qui reçoivent de la dexaméthasone.

Chez les patients recevant de l'irinotécan et de la dexaméthasone en concomitance, l'exposition au SN-38 a augmenté d'environ 110 %. Les patients doivent cesser de prendre du kétoconazole au moins une semaine avant le début du traitement par l'irinotécan. Si l'on ne peut éviter l'administration concomitante médicamenteuses avec d'autres inducteurs de la CYP3A4, il faut envisager d'augmenter la dose (voir la rubrique MISES EN GARDE ET PRÉCAUTIONS).

L'administration du CHLORHYDRATE D'IRINOTÉCAN POUR INJECTION est contre-indiquée chez les patients qui présentent une hypersensibilité à ce médicament ou à l'un des composants du produit ou du contenant.

Voir la rubrique PRÉSENTATION, COMPOSITION ET CONDITIONNEMENT pour connaître la liste complète des ingrédients.

L'administration concomitante du CHLORHYDRATE D'IRINOTÉCAN POUR INJECTION et d'inducteurs puissants de l'isoenzyme CYP3A4, tels que la phénytoïne, la carbamazépine, le phénobarbital ou la CYP3A4, est contre-indiquée puisque ces médicaments peuvent réduire la biodisponibilité et la métabolite actif SN-38 et ainsi augmenter la toxicité de l'irinotécan.

Chez les patients recevant de l'irinotécan et de la kétoconazole en concomitance, l'exposition au SN-38 a augmenté d'environ 110 %. Les patients doivent cesser de prendre du kétoconazole au moins une semaine avant le début du traitement par l'irinotécan. Si l'on ne peut éviter l'administration concomitante médicamenteuses avec d'autres inducteurs de la CYP3A4, il faut envisager d'augmenter la dose (voir la rubrique MISES EN GARDE ET PRÉCAUTIONS).

Chez les patients recevant de l'irinotécan et de la bilirubinémie en concomitance, l'exposition au SN-38 a augmenté d'environ 110 %. Les patients doivent cesser de prendre du kétoconazole au moins une semaine avant le début du traitement par l'irinotécan. Si l'on ne peut éviter l'administration concomitante médicamenteuses avec d'autres inducteurs de la CYP3A4, il faut envisager d'augmenter la dose (voir la rubrique MISES EN GARDE ET PRÉCAUTIONS).

Interactions médicament-herbe médicinale

Millepertuis : On a observé une diminution de l'exposition et du métabolite en concentrance, l'exposition au métabolite actif SN-38 est réduite d'environ 40 %. La prise de millepertuis doit être interrompue au moins 1 semaine avant le début du traitement par l'irinotécan (voir la rubrique POSOLOGIE POUR INJECTION chez la rubrique MISES EN GARDE ET PRÉCAUTIONS).

Effets du médicament sur les résultats des analyses de laboratoire

On ne connaît pas d'interaction entre l'irinotécan et les épreuves de laboratoire.

POSOLOGIE ET MODE D'ADMINISTRATION

Recommandations communes pour les schémas d'administration du CHLORHYDRATE D'IRINOTÉCAN POUR INJECTION et ajustements posologiques

Il n'existe pas de données sur l'innocuité ou l'efficacité de l'irinotécan chez les patients présentant une insuffisance hépatique importante (voir la rubrique MISES EN GARDE ET PRÉCAUTIONS).

Il faut surveiller étroitement les patients présentant des facteurs de risque d'une diarrhée grave, tels que ceux qui présentent un indice fonctionnel de 3 ou de 4 (selon la cote de l'échelle de performance du groupe d'oncologie de l'Amérique du Nord) ou ceux qui présentent une maladie