


VITAMIN B₆**PYRIDOXINE HYDROCHLORIDE
INJECTION, USP 100 mg/mL****(Vitamin B₆)**

DESCRIPTION: Pyridoxine Hydrochloride Injection (vitamin B₆) is a sterile solution of pyridoxine hydrochloride in Water for Injection with a pH between 2 and 3.8. It is intended for intramuscular or intravenous administration.

Each mL contains: Pyridoxine Hydrochloride 100 mg with Benzyl Alcohol 1.5% as preservative, in Water for Injection q.s. Sodium Hydroxide and/or Hydrochloric Acid may have been used to adjust pH.

Pyridoxine hydrochloride is a colourless or white crystal or a white crystalline powder. One gram dissolves in 5mL of Water. It is stable in air and is slowly affected by sunlight.

The structural formula of pyridoxine hydrochloride is as follows:



1. Site for - NH₂

=0

2. Site for phosphorylation

Its chemical name is 2-methyl-3-hydroxy-4, 5-bis(hydroxymethyl) pyridine hydrochloride.

CLINICAL PHARMACOLOGY: Natural substances that have vitamin B₆ activity are pyridoxine in plants and pyridoxal or pyridoxamine in animals. All 3 are converted to pyridoxal phosphate by the enzyme pyridoxal kinase. The physiologically active forms of vitamin B₆ are pyridoxal phosphate (codecarboxylase) and pyridoxamine phosphate. Riboflavin is required for the conversion of pyridoxine phosphate to pyridoxal phosphate.

Vitamin B₆ acts as a coenzyme in the metabolism of protein, carbohydrate, and fat. In protein metabolism it participates in the decarboxylation of amino acids, conversion of tryptophan to niacin or to serotonin (5-hydroxy-tryptamine), deamination, and transamination, and transulfuration of amino acids. In carbohydrate metabolism, it is responsible for the breakdown of glycogen to glucose-1-phosphate.

The total adult body pool consists of 16 to 25 mg of pyridoxine. Its half-life appears to be 15 to 20 days. Vitamin B₆ is degraded to 4-pyridoxic acid in the liver. This metabolite is excreted in the urine.

The need for pyridoxine increases with the amount of protein in the diet. The tryptophan load test appears to uncover early vitamin B₆ deficiency by detecting xanthinuria. The average adult minimum daily requirement is about 1.25 mg. The "Recommended Dietary Allowance" of the National Academy of Sciences is estimated to be as much as 2.3 for adults and 2.5 for pregnant and lactating women. The requirements are more in persons having certain genetic defects or those being treated with isonicotinic acid hydrazide (INH) or oral contraceptives.

INDICATIONS AND USAGE: Pyridoxine Hydrochloride Injection is effective for the treatment of pyridoxine deficiency as seen in the following.

Inadequate dietary intake.

Drug-induced deficiency, as from isoniazid (INH) or oral contraceptives. Inborn errors of metabolism, e.g., vitamin-B₆ dependent convulsions or vitamin-B₆ responsive anemia.

The parenteral route is indicated when oral administration is not feasible as in anorexia, nausea and vomiting, and preoperative conditions. It is also indicated when gastrointestinal adsorption is impaired.

CONTRAINDICATIONS: A history of sensitivity to pyridoxine or to any of the ingredients in the product is a contraindication.

PRECAUTION: Single deficiency, as of pyridoxine alone, is rare. Multiple vitamin deficiency is to be expected in any inadequate diet. Patients treated with levodopa should avoid supplemental vitamins that contain more than 5 mg pyridoxine in the daily dose.

Women taking oral contraceptives may exhibit increased pyridoxine requirements.

Drug Interactions: Pyridoxine supplements should not be given to patients receiving levodopa, because the action of the latter drug is antagonized by pyridoxine. However, this vitamin may be used concurrently in patients receiving a preparation containing both carbidopa and levodopa.

Use in Pregnancy-Pregnancy category A: The requirement for pyridoxine appears to be increased during pregnancy. Pyridoxine is sometimes of value in the treatment of nausea and vomiting of pregnancy.

Nursing Mothers: The need for pyridoxine is increased during lactation.

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when it is administered to a nursing woman.

Usage in Children: Safety and effectiveness in children have not been established.

ADVERSE REACTIONS: Paresthesia, somnolence, and low serum folic acid levels have been reported.

DRUG ABUSE AND DEPENDENCE: Symptoms of dependence have been noted in adults given only 200 mg daily, followed by withdrawal.

OVERDOSAGE: Pyridoxine given to animals in amounts of 3 to 4g/kg of body weight produces convulsions and death. In man, a dose of 25 mg/kg of body weight is well tolerated.

DOSAGE AND ADMINISTRATION: Pyridoxine Hydrochloride Injection may be administered by the intramuscular or intravenous route. (See INDICATIONS AND USAGE)

In cases of dietary deficiency the dosage is 10 to 20 mg daily for 3 weeks. Follow-up treatment is recommended daily for several weeks with an oral therapeutic multivitamin preparation containing 2 to 5 mg pyridoxine. Poor dietary habits should be corrected, and an adequate, well-balanced diet should be prescribed.

The vitamin B₆ dependency syndrome may require a therapeutic dosage of as much as 600 mg a day and a daily intake of 30 mg for life.

In deficiencies due to INH, the dosage is 100 mg daily for 3 weeks followed by a 30 mg maintenance dose daily.

In poisoning caused by ingestion of more than 10 grams of INH, an equal amount of pyridoxine should be given: 4 grams intravenously followed by 1 gram intramuscularly every 30 minutes.

Parenteral drug products should be inspected visually for particulate matter and discoloration whenever in solution and if the container permits.

HOW SUPPLIED: Pyridoxine Hydrochloride Injection, USP, 100 mg/mL, is available in multiple dose vials of 30 mL individually boxed.

For therapeutic use only.

Store at controlled room temperature 15° - 30° C (59° - 86° F). Do not freeze. Protect from light.

BIONICHE PHARMA (CANADA) LTD.
Belleville, ON Canada K8N 5J2

Issue Date: December 2001

0426L100